

Specific Reference Intervals of Serum Triiodothyronine, Thyroxine, and Thyroid-stimulating Hormone in Normal Pregnant Indian Women as per Trimester

¹Barun K Chakrabarty, ²Binay Mitra, ³Reetika Pal, ⁴Nandita Hazra

ABSTRACT

Aim: Maternal thyroid hormone level during pregnancy is a vital parameter for the health of mother as well as developing child. Fetal growth is affected by maternal thyroid levels. Various physiological changes like alterations of thyroxine binding globulins (TBGs), beta-human chorionic gonadotropin (β -hCG) level, and change of iodide metabolism affect maternal thyroid hormone levels. Therefore, reference intervals (RI) for thyroid hormones in pregnant population require to be established separately from general population.

Materials and methods: The RIs of serum triiodothyronine (T3), thyroxine (T4), and thyroid-stimulating hormone (TSH) were determined in healthy pregnant women by enzyme-linked immunosorbent assay (ELISA) technique after segregating them into three trimesters. This study was conducted in a 492-bedded zonal level hospital. The reference population was chosen from a study population of pregnant women by strict inclusion and exclusion criteria. The assays were done by the most commonly used economical ELISA method using standard kits. Tests were done using accurate and precise methods with proper quality control measures.

Results: The RIs were calculated from the central 95% of distribution of total T3, total T4, and TSH values located between 2.5 and 97.5 percentile values. The 0.90 confidence intervals (CIs) for the upper and lower reference limits were calculated. The values thus obtained were different from those provided by manufacturer kit literature.

Conclusion: It is recommended to determine own laboratory-specific, method-specific, trimester-wise RI for maternal thyroid hormone status and use them for screening of pregnant mothers.

Keywords: Enzyme-linked immunosorbent assay, Laboratory research, Pregnancy, Reference interval, Thyroid hormones, Trimester-specific.

How to cite this article: Chakrabarty BK, Mitra B, Pal R, Hazra N. Specific Reference Intervals of Serum Triiodothyronine, Thyroxine, and Thyroid-stimulating Hormone in Normal Pregnant Indian Women as per Trimester. *Indian J Med Biochem* 2017;21(2):96-100.

Source of support: Nil

Conflict of interest: None

INTRODUCTION

Maternal thyroid hormone level during pregnancy plays a crucial role in development of neonate and children.¹ Any gestational period thyroid hormone disorder may lead to obstetrical complications along with effects on fetus.² Fetal growth is also affected by maternal thyroid levels.³ Various physiological changes, alterations of TBGs, β -hCG level, and change of iodide metabolism affect maternal thyroid hormone levels.⁴ Therefore, to ensure optimal maternal and fetal health, it is recommended to properly assess thyroid function of all pregnant women with a distinct established reference range other than the one mentioned in the kit literature.

According to the International Federation of Clinical Chemistry (IFCC), RI is defined as an interval, i.e., bounded by two percentiles of the reference distribution, where a percentile represents a value that splits the reference distribution in a way that a stated percentage of its values has levels less than or equal to the limiting value. Most commonly, the RI of a parameter is defined as the central 95% interval bounded by the 2.5 and 97.5 percentiles of the reference distribution.⁵⁻⁸ For valid comparison to establish gestational thyroid reference range, the study population must be homogeneous to the reference individual in all respects; therefore, trimester-specific RIs of thyroid hormone can only provide the best therapeutic guideline for clinical decision making.⁹

In India and other developing and underdeveloped world, the most commonly ordered thyroid function tests are serum total T3, total T4, and TSH levels. Most common method followed is ELISA. Though free T3 and free T4 should be performed ideally as no better sensitive method is available, but due to various socioeconomic conditions, it is not yet possible for largely populated countries to conduct them.

¹Graded Specialist, ²Classified Specialist, ³Medical Officer
⁴Senior Advisor

¹Department of Pathology, Military Hospital, Ranchi, Jharkhand India

^{2,3}Department of Gynecology and Obstetrics, Military Hospital Ranchi, Jharkhand, India

⁴Department of Pathology and Microbiology, Command Hospital Lucknow, Uttar Pradesh, India

Corresponding Author: Barun K Chakrabarty, Graded Specialist Department of Pathology, Military Hospital, Ranchi, Jharkhand India, Phone: +917044648932, e-mail: bkcdc@yahoo.co.in

The purpose of our study was to determine gestational trimester-specific RIs of serum T3, T4, and TSH in Indian pregnant women in Ranchi area.

MATERIALS AND METHODS

This study was conducted in the Department of Laboratory Medicine in collaboration with the A and E Department and Family outpatient department of the Department of Gynecology and Obstetrics, of a 492-bedded hospital from June 2016 to November 2016. It is a descriptive, cross-sectional, and observational study in the reference group of pregnant women in Ranchi area.

Data were collected on the Monday of each week for a period of 06 months. A total of 240 pregnant women having single intrauterine uncomplicated pregnancy with history of consumption of iodized salt were sequentially enrolled. Data regarding gestational age were determined by last menstrual period and ultrasonography report. Case record form was filled after detailed history and examination was done by gynecologist. Written informed consent was taken.

The reference sample group was selected by applying the exclusion criteria on study population mentioned below:

- History of thyroid disorder
- History of multiple or complicated pregnancy
- History of chronic medical disorder
- Past history of spontaneous abortion
- Past or present history of antithyroid drug
- Infertility/Infertility treatment
- Regular use of any medications other than iron and folate (e.g., lithium)
- Family history of thyroid disorder
- Signs and symptoms of thyroid disorder

The remaining study population (224) created the reference sample group. The reference sample group was then segregated into groups of three trimesters according to gestational age. First trimester group constituted up to 13 weeks (182), second trimester from 13 to 27 weeks (16), and third trimester was formed by more than 28 weeks (26) gestational age ladies.

A 5 mL of fasting venous blood was collected from each participant. Collected blood serum was separated and stored at -20°C within 1 hour of collection.

Laboratory Hormonal Analysis

Serum T3, T4, and TSH assays were done using Readwell TOUCH microplate analyzer and commercially available ELISA kit namely BeneSphera™ (Avantor) by quantitative enzyme immunoassay.^{10,11} The analytical sensitivity for T3, T4, and TSH assays was 0.04 ng/mL, 0.4 µg/dL,

and 0.078 µIU/mL (1 hour incubation procedure) respectively. The intra- and inter-assay reproducibility values for T3 assays were 8.9% (for low serum range) and <5.5 and <7% (for high serum range). Similarly, for T4 these values were 6.7 and 8.3% (for low serum range) and <5% (for high serum range) and for TSH, these values are <5% (for low serum range) and <6% (for high serum range).

Statistical Analysis

All data were tabulated into MS Excel and analyzed using Statistical Package for the Social Sciences software version 19. Distribution nature of T3, T4, and TSH values of each trimester were observed by inspecting histograms. Trimester-specific mean (\bar{x}_m), standard deviation (s_x) of T3, T4, and TSH were calculated separately. The RIs were calculated by estimating the 2.5 and 97.5% determined as $\bar{X}_m \pm 1.96 * S_x$. The 0.90 CIs were obtained by the formula: Percentile $\pm 2.81 * S_x / \sqrt{n}$.⁵⁻⁸ Analysis of variance test was used to see the significance of differences among three trimester values. For TSH, nonparametric Kruskal-Wallis test was used to compare the changes between trimester values for a level of significance $p < 0.05$.

RESULTS

The study population comprised 240 pregnant women out of which 191 (79.6%) were in the first trimester, 17 (7.1%) fit into the second, and 32 (13.3%) in the third trimester. The age of the mothers ranged from 18 to 40 years. After execution of the exclusion criteria, 16 women were excluded from the study, ensuing reference sample group of 224 women, which further divided into first trimester 182 (81.3%); second trimester 16 (7.1%); and third trimester 26 (11.6%). This reference population was then used to calculate the reference limits and 0.90 trimester-specific CIs of T3, T4, and TSH.

The descriptive statistics of reference population partitioned into three trimesters are shown in Table 1. The RIs of T3, as obtained in our study, are 1.21 to 1.32 ng/mL for the first trimester; 1.13 to 1.64 ng/mL for the second trimester, and 1.16 to 1.51 ng/mL for the third trimester (Table 2). Similarly, the RIs of T4 for first, second, and third trimesters were 7.57 to 8.13, 7.17 to 8.64, and 7.07 to 8.44 µg/dL respectively, and the RIs of TSH for first, second, and third trimesters were 2.26 to 2.84, 1.61 to 3.88, and 1.59 to 2.85 µIU/mL respectively (Table 2). Box plots of T3, T4, and TSH values for the three trimesters are depicted in Graphs 1 to 3 respectively. No significant difference was observed between trimester values of all the parameters. The manufacturer provided only generalized RIs and no separate pregnancy RI level was mentioned. Generalized


Table 1: Descriptive statistics of reference population segregated into three trimesters

Trimester	Parameter	Mean ± SD	2.5 Percentile	97.5 Percentile
First trimester (n = 182)	T3 (ng/mL)	1.27 ± 0.41	1.21	1.32
	T4 (µg/dL)	7.85 ± 1.95	7.57	8.13
	TSH (µIU/mL)	2.55 ± 1.98	2.26	2.84
Second trimester (n = 16)	T3 (ng/mL)	1.39 ± 0.48	1.13	1.64
	T4 (µg/dL)	7.91 ± 1.38	7.17	8.64
	TSH (µIU/mL)	2.75 ± 2.13	1.61	3.88
Third trimester (n = 26)	T3 (ng/mL)	1.33 ± 0.08	1.16	1.51
	T4 (µg/dL)	7.76 ± 1.71	7.07	8.44
	TSH (µIU/mL)	2.22 ± 1.55	1.59	2.85


SD: Standard deviation

Table 2: Trimester-wise reference data for T3, T4, and TSH according to our study


Parameters	Trimester	Reference interval	0.90 CI of lower reference limit	0.90 CI of upper reference limit
T3 (ng/mL)	First	1.21–1.32	0.41–1.29	1.24–1.4
	Second	1.13–1.64	0.79–1.47	1.3–1.98
	Third	1.16–1.51	0.93–1.39	1.28–1.74
T4 (µg/dL)	First	7.57–8.13	7.18–7.95	7.75–8.53
	Second	7.17–8.64	6.19–8.15	7.66–9.62
	Third	7.07–8.44	6.14–7.99	7.52–9.38
TSH (µIU/mL)	First	2.26–2.84	1.84–2.68	2.42–3.26
	Second	1.61–3.88	0.12–3.09	2.4–5.36
	Third	1.59–2.85	0.75–2.43	2.01–3.69



Graph 1: Box plots for T3 values of three trimesters



Graph 2: Box plots for T4 values of three trimesters



Graph 3: Box plots for TSH values of three trimesters

RIs of T3, T4, and TSH mentioned in the kit literature were 0.52 to 1.85 ng/mL, 4.8 to 11.6 µg/dL, and 0.39 to 6.16 µIU/mL. These values are different from values obtained from our pregnant reference sample group.

DISCUSSION

Pregnancy is a crucial stage in the life of a woman. Untreated and undetected gestational thyroid disorders are known to cause many complications like anemia, pregnancy-induced hypertension, hemorrhage, preterm delivery in mothers and low birth weight, prematurity, congenital malformation, intrauterine fetal death, and lower intelligence quotient in children.¹² In recent years, various studies recommend the use of routine thyroid function tests for screening of thyroid disease during normal antenatal checkup (ANC).^{13,14} Hence, selection and proper interpretation of gestational thyroid function parameters are important for the wellbeing of both the mother and the fetus. This study is the first of its kind in Ranchi area, and presently no such pan Indian reference population data are available.

Our study provides the trimester-specific RIs of total T3, total T4, and TSH (Table 2). On comparing reference data of one trimester with the other by the t-test, it was found the change was not significant. The RI values provided by kit manufacturer were generalized and not specific for pregnancy/trimesterwise. The RIs of T3, T4, and TSH as supplied by the kit manufacturer were 0.52 to 1.85 ng/mL, 4.8 to 11.6 µg/dL, and 0.39 to 6.16 µIU/mL.

On comparing these RIs obtained and the kit manufacturer RIs, an appreciable variation of interpretation of test results was seen. This variation is because the RI provided by the reagent manufacturer has been determined on age, sex nonspecific, and nonpregnant population and cannot be applied for the interpretation of thyroid function tests of a pregnant woman.

According to the IFCC guidelines of reference values and recommendations of the Indian Thyroid Society in the line of American Thyroid Association, RI in healthy population is to be determined using stringent specific exclusion criteria.^{15,16} To fit the criteria, we have used stringent criteria for our reference population selection.

Our laboratory assessments were done by the most commonly used economical ELISA method, using standard instrument and standard kits. Tests were done using accurate and precise methods with proper quality control measures. The statistical strengths are determined by distributions examination and application of proper parametric or nonparametric methods.

Most of the previous such studies have reported trimester-specific reference ranges for free T3, free T4,

and TSH among pregnant Indian women.^{17,18} These studies reported wide variation in parameter ranges due to discrepancies in assay methods, ethnicity of study population along with variations in socioeconomic and nutritional status, reference population selection, and sample size determination criteria.^{19,20}

The strength of our study is that it is based on the population temporarily migrated to Ranchi for professional purpose and which is having pan Indian profile because of multicultural and multiethnicity and demographic distribution. The study population is considered relatively healthier than average general population resulting in representation of ideal pan Indian reference population. We have assessed total T3 and total T4 levels by ELISA method, which is most commonly assessed, economical, and most readily available parameter than free T3 and free T4 levels in our country. The limitations of our study are there is inappropriate sample size in three trimesters, second and third trimester has very small sample size as compared with first, and sample groups were not the same, i.e., the three trimesters do not have the same ANC patients, which would have provided better reference ranges. Further follow-up study with more stringent inclusion and exclusion criteria will improve our strength of data to provide more conclusive trimester-specific levels.

To conclude, it is important that each laboratory in our country should determine its own population and method-based RI for diagnosing gestational thyroid disorders. These RIs must preferentially be trimester specific, most readily available, economical, method specific, and specific for pan Indian population. To promote this concept, we have determined our own laboratory-specific, method-specific, trimester-wise RI for the most commonly used thyroid hormone parameter level by ELISA method and intend to use them for screening of Indian pregnant mothers.

REFERENCES

1. Fan X, Wu L. The impact of thyroid abnormalities during pregnancy on subsequent neuropsychological development of the offspring: a meta-analysis. *J Mater Fetal Neonatal Med* 2016 Mar;29(24):3971-3976.
2. Casey BM, Dashe JS, Wells CE, McIntire DD, Leveno KJ, Cunningham FG. Subclinical hypothyroidism and pregnancy outcomes. *Obstet Gynecol* 2005 Jul;106(1):198-199.
3. Glinoe D, Soto MF, Bourdoux P, Lejeune B, Delange F, Lemone M, Kinthaert J, Robijn C, Grun JP, de Nayer P. Pregnancy in patients with mild thyroid abnormalities: maternal and neonatal repercussions. *J Clin Endocrinol Metab* 1991 Aug;73(2):421-427.
4. Salvatore, D.; Davies, TF.; Schlumberger, MJ.; Hay, ID.; Larsen, PR. Thyroid physiology and diagnostic evaluation of patients with thyroid disorders. In: Melmed S, Polonsky K, Larsen

- PR, Kronenberg H, editors. Williams textbook of endocrinology. 12th ed. Philadelphia (PA): Saunders; 2011. pp. 327-361.
- Appold K. Determining laboratory reference intervals: CLSI guideline makes the task manageable. *Lab Med* 2009 Feb;40(2):75-76.
 - Solberg HE. Approved recommendation (1986) on the theory of reference values. Part 1. The concept of reference values. *Clin Chim Acta* 1987 May;165(1):111-118.
 - Richardson H. Medical laboratories – requirements for quality and competence: an ISO perspective. *Vox Sang* 2002 Aug;83(Suppl 1):333-335.
 - Solberg HE. Approved recommendation (1987) on the theory of reference values. Part 5. Statistical treatment of collected reference values. Determination of reference limits. *Clin Chim Acta* 1987 Dec;170(2-3):S13-S32.
 - Sahasrabudde A, Mankar J, Pitale S. Trimester specific ranges for thyroid hormones in normal pregnancy. *Thyroid Res Pract* 2016 Oct;13(3):106-109.
 - Chopra IJ, Solomon DH, Ho RS. A radioimmunoassay of thyroxine. *J Clin Endocrinol Metab* 1971 Nov;33(5):865-868.
 - Spencer CA, Takeuchi M, Kazarosyan M, MacKenzie F, Beckett GJ, Wilkinson E. Interlaboratory/intermethod differences in functional sensitivity of immunometric assays of thyrotropin (TSH) and impact on reliability of measurement of subnormal concentrations of TSH. *Clin Chem* 1995 Mar;41(3):367-374.
 - Maji R, Nath S, Lahiri S, Das MS, Bhattacharyya AR, Das HN. Establishment of trimester-specific reference intervals of serum TSH & fT4 in a pregnant Indian population at north Kolkata. *Indian J Clin Biochem* 2013 May;29(2):167-173.
 - Rajput R, Goel V, Nanda S, Rajput M, Seth S. Prevalence of thyroid dysfunction among women during the first trimester of pregnancy at a tertiary care hospital in Haryana. *Indian J Endocrinol Metab* 2015 Mar;19(3):416-419.
 - Jouyandeh Z, Hasani-Ranjbar S, Qorbani M, Larijani B. Universal screening versus selective case-based screening for thyroid disorders in pregnancy. *Endocrine* 2014 Sep;48(1):116-123.
 - Horn PS, Pesce AJ. Reference intervals: an update. *Clin Chim Acta* 2003 Aug;334(1-2):5-23.
 - Stagnaro-Green A, Abalovich M, Alexander E, Azizi F, Mestman J, Negro R, Nixon A, Pearce EN, Soldin OP, Sullivan S, et al. Guidelines of the American thyroid association for the diagnosis and management of thyroid disease during pregnancy and postpartum. *Thyroid* 2011 Oct;21(10):1081-1125.
 - Sekhri T, Juhi JA, Wilfred R, Kanwar RS, Sethi J, Bhadra K, Nair S, Singh S. Trimester specific reference intervals for thyroid function tests in normal Indian pregnant women. *Indian J Endocrinol Metab* 2016 Jan-Feb;20(1):101-107.
 - Marwaha R, Chopra S, Gopalakrishnan S, Sharma B, Kanwar RS, Sastry A, Singh S. Establishment of reference range for thyroid hormones in normal pregnant Indian women. *BJOG* 2008 Apr;115(5):602-606.
 - Jebasingh FK, Salam R, Meetei TL, Singh PT, Singh NN, Prasad L. Reference intervals in evaluation of maternal thyroid function of Manipuri women. *Indian J Endocrinol Metab* 2016 Mar-Apr;20(2):167-170.
 - Soldin OP, Soldin D, Sastoque M. Gestation-Specific thyroxine and thyroid stimulating hormone levels in the United States and worldwide. *Ther Drug Monit* 2007 Oct;29(5):553-559.